Bringing Real Life Into Every Therapy Session

Task-Specific Training, Real-World Carryover, and the Role of Immersive Therapy

42%
lower fall rate in daily life after treadmill training with a virtual reality component, compared with treadmill training alone (Mirelman et al.)

6×
lower likelihood of errors in real operations among surgical residents trained in virtual reality to an expert standard (Seymour et al.)

73% vs. 42%
of stroke survivors cleared to resume driving after simulator training versus cognitive training (Akinwuntan et al.)

Introduction

Ask most patients what they want from therapy and the answer is rarely a better score on a standardized test. They want to button a shirt before work, carry groceries in from the car, step off a curb without hesitating, or drive themselves to a family event. Clinicians know this, which is why so much of modern rehabilitation is built around task-specific training, the practice of goal-directed, functional movements that resemble the activities a person needs to perform. The approach rests on a well-established principle of neuroplasticity. Kleim and Jones list specificity among the core principles of experience-dependent plasticity, explaining that the nature of the training experience shapes the nature of the change that follows.

Specificity explains why practice should look like the real task. The next question, and the one patients care about most, is whether a skill practiced in therapy carries over into the kitchen, the grocery store, or the car. Rehabilitation researchers call this carryover or transfer, and it corresponds to what the World Health Organization's International Classification of Functioning, Disability and Health describes as performance, meaning what a person actually does in their everyday environment rather than what they can do under test conditions (World Health Organization). This paper reviews research showing that skills transfer best when practice resembles the real context of a task, and that simulated and virtual environments have produced gains that appear in real-world outcomes such as safer walking, return to driving, and everyday memory. The final sections connect these findings to the design of the Smart Therapy Complete Solution, framed as design rationale drawn from the research rather than as outcome claims about the product itself.

Why the Practice Context Matters

Learning research has long recognized that skills travel best over short distances. Barnett and Ceci proposed a taxonomy of transfer that separates the content of what is learned from the context in which it is used, including the physical setting, the functional purpose, the social situation, and the modality of the task. Transfer is most dependable when these contexts match, and each difference between the practice setting and the real one adds to the distance a skill must cover. A reach practiced at a clinic table has a long way to go before it becomes a reach into a kitchen cabinet while dinner is on the stove, whereas a reach practiced inside a realistic kitchen scene starts much closer to that goal.

Kleim and Jones describe a similar idea at the level of the brain. Their principle of specificity holds that the nature of the training experience shapes the nature of the resulting plasticity, and their principle of transference holds that plasticity from one training experience can enhance the acquisition of similar behaviors. Together, these principles suggest that practice designed to resemble the target activity in its movements, goals, and surroundings gives the nervous system the best opportunity to apply what was learned when the real activity comes along.

The structure of practice matters as well. In a classic experiment, Shea and Morgan had participants learn movement patterns either in blocks, repeating one pattern many times before moving on, or in a random order. Random practice was more demanding during acquisition but produced better retention and transfer. Because real life rarely presents the same task twice in a row, practice that varies its demands tends to prepare learners better for the world outside the clinic.

These principles create a practical challenge for conventional therapy, since a clinic gym can only partly reproduce the visual, spatial, and cognitive demands of a home, a store, or a busy street, and changing those demands often means rearranging equipment. Levac, Huber, and Sternad argue that virtual environments answer this challenge by offering flexible, realistic platforms for practicing complex skills without giving up control over the conditions of practice, and they conclude that virtual environments are well positioned to enable transfer from therapeutic practice to the real world. They also point out that the fidelity and dimensionality of a virtual environment shape how skills are learned and transferred, which is one reason three-dimensional, immersive presentations of everyday tasks have drawn growing interest in rehabilitation.

Evidence That Simulated Practice Carries Into Real Life

A growing body of trials has measured outcomes in the real world after training in simulated or virtual environments. The findings come from surgery, driving, fall prevention, and everyday cognition, and together they show that practice in a well-designed simulation can change what people do outside of it.

Surgical Skill


Some of the most direct evidence comes from surgical education. In a randomized, double-blinded study, Seymour and colleagues trained eight surgical residents on a

virtual reality simulator until each reached criterion levels set by experienced laparoscopic surgeons, while eight comparison residents received no simulator training. During real gallbladder procedures scored by blinded reviewers, the VR-trained residents performed the dissection 29 percent faster, and errors were six times less likely (1.19 versus 7.38 errors per case). Two features of this design carry over well to rehabilitation, since training was tied to a performance standard rather than a fixed amount of time and the virtual task closely matched the real one.

Returning to the Road After Stroke

Driving rehabilitation offers a parallel example. Akinwuntan and colleagues randomized 83 people with first-ever subacute stroke to 15 hours of simulator-based driving training or 15 hours of driving-related cognitive tasks over five weeks. Both groups improved after training, but the simulator group's advantage appeared where it mattered most, since 73 percent of the simulator group passed the official predriving assessment six to nine months after stroke and were legally allowed to resume driving, compared with 42 percent of the cognitive training group. A further analysis of the same trial found that simulator-trained participants scored better on the overall on-road test and on specific items including anticipation and perception of signs, visual behavior and communication, quality of traffic participation, and turning left (Devos et al.).

Figure 1. Real-world outcomes after simulated practice




Left: share of stroke survivors who passed the official predriving assessment six to nine months after stroke (n = 83). Right: mean errors per case during real laparoscopic gallbladder procedures (n = 16). Sources: Akinwuntan et al., Neurology, 2005; Seymour et al., Annals of Surgery, 2002.

Walking Safely and Preventing Falls

Fall prevention research has produced one of the largest real-world trials of virtual training. Mirelman and colleagues tested whether adding a virtual reality component to treadmill training would reduce falls in daily life among 282 older adults aged 60 to 90

who were at high risk of falling, a group that included people with Parkinson's disease and mild cognitive impairment. In the virtual reality group, a camera projected participants' feet onto a screen in front of the treadmill, where they navigated real-life challenges such as stepping over obstacles and puddles and following pathways. In the six months after training, the fall rate in the virtual reality group dropped significantly from its pretraining level, while the decrease in the treadmill-only group was not significant. Compared directly, the virtual reality group's fall rate was 42 percent lower than that of the treadmill-only group (incident rate ratio 0.58, 95% CI 0.36 to 0.96), and no serious training-related adverse events occurred.

Figure 2. Falls in daily life before and after training



Falls per six months before and after treadmill training with or without a virtual reality component (N = 282 older adults at high fall risk). Source: Mirelman et al., The Lancet, 2016.

The V-TIME system was screen-based rather than fully immersive, which makes the result especially informative, because it suggests that bringing realistic environmental and cognitive challenges into walking practice is itself an active ingredient. Other fall prevention research points the same way. In the LiFE trial, Clemson and colleagues built balance and strength training into older adults' daily routines and reduced the rate of falls by 31 percent compared with a gentle exercise control (incidence rate ratio 0.69, 95% CI 0.48 to 0.99), while a structured program of similar exercises performed three times a week did not produce a statistically significant reduction. Training that brings real-world demands into practice appears to protect people better in the real world.

Everyday Memory and Thinking

Virtual environments built around daily activities have also produced real-life gains in cognition. Yip and Man developed a 12-session virtual reality program for people with acquired brain injury that used everyday prospective memory activities, such as remembering to carry out an intended task at the right moment, as its training content.


Participants showed significantly better changes on both virtual and real-life prospective memory measures, along with improvements in frontal lobe function and semantic fluency.

Faria and colleagues compared Reh@City v2.0, an adaptive program that trains cognition through simulations of everyday tasks in a virtual city, with paper-and-pencil training of equivalent content in 36 people with chronic stroke. The virtual city group showed greater improvements in general cognitive functioning, visuospatial ability, and executive functions, and its gains extended to self-perceived cognitive difficulties in everyday life. The authors concluded that the intervention with higher ecological validity was the more effective of the two.

Why Rehabilitation-Specific and Immersive Design Matters

Within stroke upper limb rehabilitation, the evidence favors virtual reality that is built for therapy. Maier and colleagues pooled 30 randomized trials involving 1,473 patients and separated systems designed specifically for rehabilitation from recreational, off-the-shelf gaming systems. Compared with conventional therapy, rehabilitation-specific systems showed significant effects on body function (SMD 0.23, 95% CI 0.10 to 0.36) and activity (SMD 0.31, 95% CI 0.15 to 0.47), with moderate certainty of evidence, while recreational systems did not show a significant effect on either outcome. The authors identified six principles present in more than half of the rehabilitation-specific systems: variable practice, promoting use of the affected limb, implicit feedback about movement quality, increasing difficulty, task-specific practice, and explicit feedback about results.

Figure 3. Rehabilitation-specific versus recreational VR for upper limb recovery after stroke



Pooled standardized mean differences compared with conventional therapy across 30 randomized trials (1,473 patients). Confidence intervals that cross zero indicate no significant effect. Source: Maier et al., Neurorehabilitation and Neural Repair, 2019.

Fully immersive, head-mounted systems are the newest part of this literature, and early results are encouraging. Kiper and colleagues identified 10 randomized trials with 324 participants comparing immersive VR with conventional rehabilitation after stroke. Immersive VR was associated with better upper extremity function on the Fugl-Meyer Assessment (mean difference 6.33 points, 95% CI 4.15 to 8.50) and better performance of activities of daily living (SMD 0.58, 95% CI 0.25 to 0.91). Several of the included programs built practice around everyday tasks such as virtual cooking, tidying a desk, and household chores. Because the trials were recent and generally small, the authors called for larger studies with long-term follow-up to confirm how these gains carry into daily life.

Strengthening the Bridge to Daily Life

Simulation narrows the distance between practice and real life, and carryover research shows how a few deliberate steps can close the remaining gap. Taub and colleagues added a transfer package to constraint-induced movement therapy for 40 adults at least one year after stroke. The package included behavioral contracts, a daily home diary, structured review of arm use in everyday activities, problem solving around barriers, home practice assignments, and weekly phone calls after treatment. With the same amount of in-clinic time, participants who received the package showed gains in real-world arm use that were 2.4 times as large as those without it, and the gains were retained one year later (Taub et al.).

Teaching patients to solve problems for themselves extends transfer even further. In an exploratory randomized trial, McEwen and colleagues combined task-specific practice with a global problem-solving strategy known as Goal, Plan, Do, Check, an approach called Cognitive Orientation to daily Occupational Performance (CO-OP). Compared with usual care, CO-OP produced a large effect on activities participants had never practiced in therapy (d = 1.2 immediately after treatment and d = 1.1 at three months), as well as a large effect on trained activities at follow-up (d = 1.6).

Table 1 summarizes the studies in this paper that measured real-world or everyday outcomes after simulated or virtual practice.

Table 1. Studies showing everyday gains after simulated or virtual practice

Study	Who was studied	Simulated practice	Real-world outcome	Key finding
Seymour et al. (2002)	16 surgical residents	VR simulator training to an expert criterion	Errors during real surgery	29 percent faster dissection and errors six times less likely
Akinwuntan et al. (2005)	83 adults with subacute stroke	15 hours of driving simulator training	Official predriving assessment	73 percent vs. 42 percent cleared to resume driving
Mirelman et al. (2016)	282 older adults at high fall risk	Treadmill walking with virtual obstacles	Falls in daily life over six months	42 percent lower fall rate than treadmill training alone
Yip and Man (2013)	Adults with acquired brain injury	12 sessions of everyday memory tasks in VR	Real-life prospective memory tasks	Significantly better changes on real-life memory measures
Faria et al. (2020)	36 adults with chronic stroke	Daily tasks in a virtual city	Cognition and self-perceived everyday deficits	Greater gains than equivalent paper-and-pencil training
Kiper et al. (2024)	10 trials, 324 adults after stroke	Immersive head-mounted VR	Activities of daily living	Moderate effect favoring immersive VR (SMD 0.58)

Designing Immersive Sessions With Carryover in Mind

The Smart Therapy Complete Solution pairs a Pico Neo 3 Pro headset worn by the patient with a tablet the therapist uses to control the experience, see what the patient sees in real time, and adjust exercise settings until the task is challenging but still achievable. Exercises support physical, occupational, and neurologic therapy, and each tablet can control any headset assigned to the clinic. Table 2 connects the carryover principles in this paper to ways a therapist can apply them during a session. These connections describe design intent drawn from published findings, and they are not claims that the product itself has been shown to improve real-world carryover.

Table 2. Carryover principles and how they can look in a Smart Therapy session

Carryover principle	Research basis	In a Smart Therapy session
Match practice to the real task	Specificity and context matching support transfer, and closely matched simulations transferred in surgery, driving, and everyday cognition (Kleim and Jones; Barnett and Ceci; Seymour et al.; Faria et al.)	**Plan:** choose exercises whose movements and goals line up with the daily activities the patient has named as priorities.
Bring real-world challenge into practice	Walking practice with virtual obstacles reduced falls in daily life by 42 percent (Mirelman et al.)	**Deliver:** increase visual, spatial, and cognitive demands as skill grows, using the live view to keep the challenge achievable.
Vary the conditions of practice	Random practice improved retention and transfer, and variable practice appeared in 86 percent of rehabilitation-specific VR trials (Shea and Morgan; Maier et al.)	**Deliver:** change exercise settings within and across sessions so the patient meets the task in more than one form.
Coach strategy along with movement	Cognitive strategy training produced large effects on untrained activities (McEwen et al.)	**Deliver:** use the live tablet view to ask the patient what happened and how they might adjust, then let them test their own solution.
Build the bridge to daily life	A transfer package produced 2.4 times larger real-world gains (Taub et al.), and training built into routines reduced falls (Clemson et al.)	**Document:** record the exercises and settings used so home assignments and follow-up conversations connect directly to what was practiced.

The common thread is that the headset session becomes a realistic rehearsal space within a progression that ends in the patient's own environment. Immersive practice lets a patient meet the visual and spatial demands of a task in a controlled setting, and the therapist's live view makes it possible to adjust difficulty and coach strategy as practice unfolds. Pairing those sessions with real-task practice in the clinic, specific activities at home, and follow-up on how the skill is being used reflects the approaches with the strongest carryover evidence.

Considerations for Clinics

For clinics evaluating immersive tools with carryover in mind, the research suggests a few practical questions. The first is whether the system was designed for rehabilitation, since rehabilitation-specific VR has shown clearer benefits than recreational gaming systems (Maier et al.). The second is whether exercises resemble the daily activities patients care about and can be varied as skill grows. The third is whether the therapist keeps control over difficulty and coaching throughout the session. The fourth is whether gains are tracked outside the clinic, for example with the Motor Activity Log, a

structured interview about everyday arm use that correlates strongly with accelerometry (r = 0.71 to 0.91), so the team can see progress reaching daily life (Taub et al.).

Patient selection and tolerance also deserve attention. Immersive VR trials have generally reported good acceptability, but some participants have experienced side effects such as eye strain, blurred vision, sweating, headache, dizziness, or nausea (Kiper et al.), so screening for tolerance and observing patients during immersive exercises remain part of good practice with any headset-based system.

Conclusion

Task-specific training remains one of the best-supported principles in rehabilitation, and the research reviewed here shows how to extend its reach. Skills transfer best when practice resembles the real context of a task, and trials in surgery, driving, fall prevention, and everyday cognition show that well-designed simulated practice can produce gains people carry into real life. Rehabilitation-specific and immersive systems add realistic, three-dimensional versions of everyday tasks to that toolkit, with encouraging early results for function and activities of daily living.

Within the Plan, Deliver, Measure, Document workflow of the Smart Therapy Complete Solution, the therapist plans exercises around the patient's daily-life goals, delivers varied and appropriately challenging practice with a live view of the patient's experience, measures progress from one visit to the next, and documents the progression that links each session to the next step outside the clinic. The research suggests that this combination of realistic practice and a deliberate bridge to daily life.

Works Cited

- Akinwuntan AE, PhD, et al. *Effect of Simulator Training on Driving After Stroke: A Randomized Controlled Trial*. www.nads-sc.uiowa.edu/dscna/2005/papers/Effect_of_simulator_training_driving_after_stroke.pdf. Accessed 20 Oct. 2026.
- Barnett, Susan M., and Stephen J. Ceci. “When and Where Do We Apply What We Learn?: A Taxonomy for Far Transfer..” *Psychological Bulletin*, vol. 128, no. 4, 2002, pp. 612–637, psycnet.apa.org/doi/10.1037/0033-2909.128.4.612, 10.1037/0033-2909.128.4.612. Accessed 20 Oct. 2026.
- Clemson, L., et al. “Integration of Balance and Strength Training Into Daily Life Activity to Reduce Rate of Falls in Older People (the LiFE Study): Randomised Parallel Trial.” *BMJ*, vol. 345, no. aug07 1, 7 Aug. 2012, pp. e4547-e4547, www.bmj.com/content/345/bmj.e4547, 10.1136/bmj.e4547. Accessed 20 Oct. 2026.
- Devos, Hannes, et al., 19386794. “Comparison of the Effect of Two Driving Retraining Programs on On-Road Performance After Stroke.” *Neurorehabilitation and Neural Repair*, vol. 23, no. 7, Sept. 2009, pp. 699–705. *PubMed*, pubmed.ncbi.nlm.nih.gov/19386794/, 10.1177/1545968309334208. Accessed 20 Oct. 2026.
- Faria, Ana Lúcia, et al. “A Comparison of Two Personalization and Adaptive Cognitive Rehabilitation Approaches: A Randomized Controlled Trial With Chronic Stroke Patients.” *Journal of NeuroEngineering and Rehabilitation*, vol. 17, no. 1, 16 June 2020, 10.1186/s12984-020-00691-5.
- Kiper, Pawel, et al. “Effects of Immersive Virtual Reality on Upper-Extremity Stroke Rehabilitation: A Systematic Review With Meta-Analysis.” *Journal of Clinical Medicine*, vol. 13, no. 1, 1 Jan. 2024, p. 146, www.mdpi.com/2077-0383/13/1/146#:~:text=The%20results%20show%20that%20immersive, 10.3390/jcm13010146. Accessed 20 Oct. 2026.

- Kleim, Jeffrey A., and Theresa A. Jones. “Principles of Experience-Dependent Neural Plasticity: Implications for Rehabilitation After Brain Damage.” *Journal of Speech, Language, and Hearing Research*, vol. 51, no. 1, 2008, https://doi.org/10.1044/1092-4388(2008/018).
- Levac, Danielle E., et al. “Learning and Transfer of Complex Motor Skills in Virtual Reality: A Perspective Review.” *Journal of NeuroEngineering and Rehabilitation*, vol. 16, no. 1, 18 Oct. 2019, link.springer.com/article/10.1186/s12984-019-0587-8, 10.1186/s12984-019-0587-8. Accessed 20 Oct. 2026.
- Maier, Martina, et al. “Effect of Specific Over Nonspecific VR-Based Rehabilitation on Poststroke Motor Recovery: A Systematic Meta-Analysis.” *Neurorehabilitation and Neural Repair*, vol. 33, no. 2, 1 Feb. 2019, pp. 112–129, www.ncbi.nlm.nih.gov/pubmed/30700224, 10.1177/1545968318820169. Accessed 20 Oct. 2026.
- McEwen, Sara, et al. “Combined Cognitive-Strategy and Task-Specific Training Improve Transfer to Untrained Activities in Subacute Stroke.” *Neurorehabilitation and Neural Repair*, vol. 29, no. 6, 21 Nov. 2014, pp. 526–536, 10.1177/1545968314558602.
- Mirelman, Anat, et al. “Addition of a Non-Immersive Virtual Reality Component to Treadmill Training to Reduce Fall Risk in Older Adults (V-TIME): A Randomised Controlled Trial.” *The Lancet*, vol. 388, no. 10050, Sept. 2016, pp. 1170–1182, 10.1016/s0140-6736(16)31325-3.
- Seymour, Neal E, et al. “Virtual Reality Training Improves Operating Room Performance: Results of a Randomized, Double-Blinded Study.” *Annals of Surgery*, vol. 236, no. 4, 2002, pp. 458–63; discussion 463–4, www.ncbi.nlm.nih.gov/pmc/articles/PMC1422600/., 10.1097/00000658-200210000-00008. Accessed 20 Oct. 2026.
- Shea, John B, and Robyn L Morgan. “(PDF) Contextual Interference Effects on the Acquisition, Retention, and Transfer of a Motor Skill.” *ResearchGate*, Mar. 1979, www.researchgate.net/publication/263935183_Contextual_interference_effects_on_the_acquisition_retention_and_transfer_of_a_motor_skill. Accessed 20 Oct. 2026.
- Taub, Edward, et al. “Method for Enhancing Real-World Use of a More Affected Arm in Chronic Stroke.” *Stroke*, vol. 44, no. 5, May 2013, pp. 1383–1388, 10.1161/strokeaha.111.000559.
- World Health Organisation. “International Classification of Functioning, Disability and Health (ICF).” *World Health Organisation*, 2001, www.who.int/standards/classifications/international-classification-of-functioning-disability-and-health. Accessed 20 Oct. 2026.
- Yip, Ben C B, and David W K Man, 23422463. “Virtual Reality-Based Prospective Memory Training Program for People With Acquired Brain Injury.” *NeuroRehabilitation*, vol. 32, no. 1, 2013, pp. 103–15. *PubMed*, pubmed.ncbi.nlm.nih.gov/23422463/, 10.3233/NRE-130827. Accessed 20 Oct. 2026.